arXiv:math/0401431v1 [math.AT] 30 Jan 2004

COINCIDENCE CLASSES IN NONORIENTABLE MANIFOLDS

DANIEL VENDRÚSCOLO

Abstract. In this article we studied Nielsen coincidence theory for maps between manifolds of same dimension without hypotheses on orientation. We use the definition of semi-index of a class, we review the definition of defective classes and study the appearance of defective root classes. We proof a semi-index product formula type for lifting maps and we presented conditions such that defective coincidence classes are the only essencial classes.

1. Introduction

Nielsen coincidence theory was extended ([Dobreńko & Jezierski] and [Jezierski 2]) to maps between nonorientable topological manifolds using the notion of $semi-index$ (a non negative integer) for a coincidence set.

We consider maps $f, g : M \to N$ between manifolds without boundary of the same dimension n, we define $h = (f, g) : M \to N \times N$, then using microbundles (see [Jezierski 2] for details) we can suppose that h is in a transverse position.

Let w be a path satisfying the Nielsen relation between $x, y \in Coin(f, g)$. We choose a local orientation γ_0 of M in x and denote by γ_t the translation of γ_0 along $w(t)$.

Definition 1.1. [Jezierski 2, 1.2] *We will say that two points* $x, y \in Coin(f, g)$ *are R-related* (xRy) *if and only if there is a path* w *establishing the Nielsen relation between them such that the translation of the orientation* $h_*\gamma_0$ *along a path in the diagonal* $\Delta(N) \subset N \times N$ *homotopic to* $h \circ w$ *in* $N \times N$ *is opposite to* $h_*\gamma_1$*. In this case the path* w *is called graph-orientation-reversing.*

Since (f, g) is transverse, $Coin(f, g)$ is finite. Let $A \subset Coin(f, g)$, then A can be represented as $A = \{a_1, a_2, \cdots, a_s; b_1, c_1, \cdots, b_k, c_k\}$ where b_iRc_i for any i and a_iRa_j for no $i \neq j$. The elements $\{a_i\}_i$ of this decomposition are called *free.*

Definition 1.2. *In the above situation the* $semi - index$ *of the pair* (f, g) *in* $A = \{a_1, a_2, \cdots, a_s; b_1, c_1, \cdots, b_k, c_k\}$ *(denoted* $|ind|(f, g; A)$*)*

2000 *Mathematics Subject Classification.* Primary 55M20.

Key words and phrases. Nielsen theory, coincidence theory, coincidence index.

is the number of free elements of this decomposition of A.

$$|ind|(f,g;A)=s.$$

In [Dobreńko & Jezierski] and [Jezierski 2] we can find the proof of the fact that this definition does not depend on the decomposition of A. Moreover, if $U\subset M$ is an open subset we can extend this definition to a semi-index of a pair on the subset U ($|ind|(f,g;U)$).

Definition 1.3. *A coincidence class* C *of a transverse pair* (f,g) *is essential if* $|ind|(f,g;A)\neq 0$.

In [Jezierski 1] the authors studied when a coincidence point $x\in Coin(f,g)$ satisfy xRx. These points can only appear in the nonorientable case, they will be called *self-reducing* points. They represent a new situation (see [Jezierski 1, Example 2.4]) that never occurs in the orientable case or in the fixed point context.

Definition 1.4. *(*[Jezierski 1, 2.1]*) Let* $x\in Coin(f,g)$ *and let* $H\subset\pi_1(M)$, $H'\subset\pi_1(N)$ *denote the subgroups of orientation-preserving elements. We define:*

$$Coin(f_\#,g_\#)_x=\{\alpha\in\pi_1(M,x)\mid f_\#(\alpha)=g_\#(\alpha)\},$$
$$Coin^+(f_\#,g_\#)_x=Coin(f_\#,g_\#)_x\cap H.$$

Lemma 1.5. *(*[Jezierski 1, 2.2]*) Let* $f,g:M\to N$ *be transverse and* $x\in Coin(f,g)$. *Then* xRx *if and only if* $Coin^+(f_\#,g_\#)_x\neq Coin(f_\#,g_\#)_x\cap f_\#^{-1}(H')$ *(in other words, if there exists a loop* α *based at* x *such that* $f\circ\alpha\sim g\circ\alpha$ *and exactly one of the loops* α *or* $f\circ\alpha$ *is orientation-preserving).*

Definition 1.6. *A coincidence class* A *is called defective if* A *contains a self-reducing point.*

Lemma 1.7. *(*[Jezierski 1, 2.3]*) If a Nielsen class* A *contains a self-reducing point (i.e.* A *is defective) then any two points in this class are* R*-related, and thus:*

$$|ind|(f,g;A)=\begin{cases}0 & \text{if } \#A \text{ is even;}\\ 1 & \text{if } \#A \text{ is odd.}\end{cases}$$

2. The root case

In [Brown & Schirmer] we can find a different approach to extending the Nielsen root theory to the nonorientable case. Using the concept of *orientation-true*[1] map they classified maps between manifolds of the same dimension in three types (see also [Olum] and [Skora]):

Definition 2.1. *(*[Brown & Schirmer, 2.1]*) Let* $f:M\to N$ *be a map of manifolds. Then three types of maps are defined as follows.*

---

[1] f is orientation-true if for each loop $\alpha\in\pi_1(M)$, $f(\alpha)$ is orientation-preserving if and only if α is orientation-preserving.

(1) *Type I: f is orientation-true.*
(2) *Type II: f is not orientation-true but does not map an orientation-reversing loop in M to a contractible loop in N.*
(3) *Type III: f maps an orientation-reversing loop in M to a contractible loop in N.*

Further, a map f is defined to be orientable if it is of Type I or II, and nonorientable otherwise.

For orientable maps they describe an *Orientation Procedure* ([Brown & Schirmer, 2.6]) for root classes, using local degree with coefficients in $\mathbb{Z}$. For maps of Type III the same procedure is only possible with coefficients in $\mathbb{Z}_2$. They then defined the *multiplicity* of a root class, that is an integer for orientable maps and an element of $\mathbb{Z}_2$ for maps of Type III

Now if we consider the root classes for a map f to be the coincidence classes of the pair (f, c) where c is the constant map, we have:

Theorem 2.2. *Let $f : M \to N$ be a map of manifolds of the same dimension.*

(i) *If f is orientable, no root class of f is defective.*
(ii) *If f is of Type III, then all root classes of f are defective.*

Proof: By lemma 1.5, a coincidence class C of the pair (f, c) is defective if and only if there exists a point $x \in C$ and a loop α in x such that $f \circ \alpha \sim 1$ and α is orientation-reversing. This proves the first statement.

Now let f be a Type III map. Then there exists a loop $\alpha \in \pi_1(M, x_0)$ such that α is orientation-reversing and $f \circ \alpha \sim 1$. If x is a root of f choosing a path β from x to x_0 we have that $\gamma = \beta^{-1}\alpha\beta$ is a loop in x such that γ is orientation-reversing and $f \circ \gamma \sim 1$. Then all roots of f are self-reducing points. □

In fact [Brown & Schirmer, Lemma 4.1] shows the equality between the multiplicity of a root class and its semi-index.

Theorem 2.3. *Let M and N be manifolds of the same dimension such that M is nonorientable and N is orientable. If $f : M \to N$ is a map then all essential root classes of f are defective.*

Proof: There are no orientation-true maps from a nonorientable to an orientable manifold. If f is a Type II map then by [Brown & Schirmer, Lemma 3.10] $deg(f) = 0$ and f has no essential root classes. The result follows by proposition 2.2. □

Using the ideas of Proposition 2.2 we can also state:

Lemma 2.4. *Let $f, g : M \to N$ be two maps between manifolds of the same dimension. If there exist a coincidence point x_0 and a graph-orientation-reverse loop α in x_0 such that $f(\alpha)$ is in the center of $\pi_1(N)$, then all coincidence points of the pair (f, g) are self-reducing points.*

Proof: We can suppose that all coincidences of the pair (f, g) have image on the same point in N. If $x_1 \in Coin(f, g)$ we choose a path β from x_1 to x_0 and take the loop $\gamma = \beta^{-1} \circ \alpha \circ \beta$ at x_1. Since α belongs to the center of $\pi_1(N)$, γ is a graph-orientation-reverse loop at x_1. $\square$

Corollary 2.5. *Let $f, g : M \to N$ be two maps between manifolds of the same dimension such that $f_\#(\pi_1(M))$ is contained in the center of $\pi_1(N)$. If (f, g) has a defective class then all classes of (f, g) are defective.*

3. COVERING MAPS

Let M and N be compact, closed manifolds of the same dimension; $f, g : M \to N$ be two maps such that $Coin(f, g)$ is finite. and $p : \widetilde{M} \to M$ and $q : \widetilde{N} \to N$ be finite coverings such that there exist lifts $\widetilde{f}, \widetilde{g} : \widetilde{M} \to \widetilde{N}$ of the pair f, g.

$$\begin{tikzcd}
\widetilde{M} \arrow[r, "\widetilde{f}", "\widetilde{g}"'] \arrow[d, "p"'] & \widetilde{N} \arrow[d, "q"] \\
M \arrow[r, "f", "g"'] & N
\end{tikzcd}$$

In this situation the groups of Deck transformations of the lifts $\widetilde{M}$ and $\widetilde{N}$ can be described by:

$$\mathfrak{D}(\widetilde{M}) = \frac{\pi_1(M)}{p_\#(\pi_1(\widetilde{M}))}; \quad \mathfrak{D}(\widetilde{N}) = \frac{\pi_1(N)}{q_\#(\pi_1(\widetilde{N}))}$$

Choosing a point $x_0 \in Coin(f, g)$ we define $\widetilde{f}_{*,x_0}, \widetilde{g}_{*,x_0} : \mathfrak{D}(\widetilde{M}) \to \mathfrak{D}(\widetilde{N})$ such that, for each $\alpha \in \mathfrak{D}(\widetilde{M})$ we take $\overline{\alpha} \in \pi_1(M, x_0)$ such that $\mathfrak{D}(\widetilde{M}) \ni [\overline{\alpha}] = \alpha$ and:

$$\widetilde{f}_{*,x_0}(\alpha) = [f_\#(\overline{\alpha})] \in \mathfrak{D}(\widetilde{N})$$

$$\widetilde{g}_{*,x_0}(\alpha) = [g_\#(\overline{\alpha})] \in \mathfrak{D}(\widetilde{N})$$

We note that $\widetilde{f}_{*,x_0}, \widetilde{g}_{*,x_0}$ may depend on x_0. Choosing a $x_0 \in Coin(f, g)$ we have. $\forall \widetilde{x} \in \widetilde{M}$ and $\forall \alpha \in \mathfrak{D}(\widetilde{M})$:

$$\widetilde{f}(\alpha(\widetilde{x})) = \widetilde{f}_{*,x_0}(\alpha) \circ \widetilde{f}(\widetilde{x})$$

$$\widetilde{g}(\alpha(\widetilde{x})) = \widetilde{g}_{*,x_0}(\alpha) \circ \widetilde{g}(\widetilde{x})$$

Lemma 3.1. *Let $\widetilde{x_0} \in Coin(\widetilde{f}, \widetilde{g})$ and $\alpha \in \mathfrak{D}(\widetilde{M})$. Then $\alpha(\widetilde{x_0}) \in Coin(\widetilde{f}, \widetilde{g})$ if and only if $\widetilde{f}_{*,x_0}(\alpha) = \widetilde{g}_{*,x_0}(\alpha)$ where $x_0 = p(\widetilde{x_0})$.*

Proof: By the above observation we have $\widetilde{f}(\alpha(\widetilde{x_0})) = \widetilde{f}_{*,x_0}(\alpha) \circ \widetilde{f}(\widetilde{x_0})$ and $\widetilde{g}(\alpha(\widetilde{x_0})) = \widetilde{g}_{*,x_0}(\alpha) \circ \widetilde{g}(\widetilde{x_0})$. This proves the lemma. $\square$

Corollary 3.2. *Let* $\widetilde{x_0} \in Coin(\widetilde{f},\widetilde{g})$ *and* $x_0 = p(\widetilde{x_0})$*. Then* $p^{-1}(x_0) \cap Coin(\widetilde{f},\widetilde{g})$ *have exactly* $\#Coin(\widetilde{f}_{*,x_0},\widetilde{g}_{*,x_0})$ *elements.* □

Lemma 3.3. *Let* $\widetilde{x_0}$ *and* $\widetilde{x}'_0$ *be two coincidences of the pair* $(\widetilde{f},\widetilde{g})$ *such that* $p(\widetilde{x_0}) = p(\widetilde{x}'_0) = x_0$*, and let* γ *the unique element of* $\mathfrak{D}(\widetilde{M})$ *such that* $\gamma(\widetilde{x_0}) = \widetilde{x}'_0$*. The points* $\widetilde{x_0}$ *and* $\widetilde{x}'_0$ *are in the same coincidence class of* $(\widetilde{f},\widetilde{g})$ *if and only if there exists* $\overline{\gamma} \in \pi_1(M,x_0)$ *such that:*

- $\mathfrak{D}(\widetilde{M}) \ni [\overline{\gamma}] = \gamma$;
- $f_\#(\overline{\gamma}) = g_\#(\overline{\gamma})$.

Proof: $(\Rightarrow)$ If $\widetilde{x_0}$ and $\widetilde{x}'_0$ are in the same coincidence class of $(\widetilde{f},\widetilde{g})$, there exists a path β from $\widetilde{x_0}$ to $\widetilde{x}'_0$ that realizes the Nielsen relation, (*i.e.* $\widetilde{f}\circ\beta \sim \widetilde{g}\circ\beta$).

Take $\overline{\gamma} = p(\beta) \in \pi_1(M,x_0)$. We can see that $[\overline{\gamma}] = \gamma$ and $f\circ\overline{\gamma} = q\circ\widetilde{f}\circ\beta \sim q\circ\widetilde{g}\circ\beta = g\circ\overline{\gamma}$ showing that $f_\#(\overline{\gamma}) = g_\#(\overline{\gamma})$.

$(\Leftarrow)$ Take the lift $\widetilde{\gamma}$ of $\overline{\gamma}$ starting at $\widetilde{x_0}$. It is a path from $\widetilde{x_0}$ to $\widetilde{x}'_0$ that realizes the Nielsen relation, (*i.e.* $\widetilde{f}\circ\widetilde{\gamma} \sim \widetilde{g}\circ\widetilde{\gamma}$). □

Corollary 3.4. *In lemma 3.3, if the points* $\widetilde{x_0}$ *and* $\widetilde{x}'_0$ *are in the same coincidence class of* $(\widetilde{f},\widetilde{g})$ *then* $\widetilde{x_0}R\widetilde{x}'_0$ *if and only if* $sign(\widetilde{f}_{*,x_0}(\gamma))\cdot sign(\gamma) = -1$*. In this case,* x_0 *is a self-reducing coincidence point.*

Proof: First we note that since $f_\#(\overline{\gamma}) = g_\#(\overline{\gamma})$ then $\widetilde{f}_{*,x_0}(\gamma) = \widetilde{g}_{*,x_0}(\gamma)$ and we can see that $sign(\widetilde{f}_{*,x_0}(\gamma))\cdot sign(\gamma) = -1$ if and only if the paths $\overline{\gamma}$ and $\widetilde{\gamma}$ in the proof of lemma 3.3 are both graph orientation-reversing. □

Denoting the natural projection by $j_{x_0} : \pi_1(M,x_0) \to \mathfrak{D}(\widetilde{M})$ and the set $\{\overline{\alpha} \in \pi_1(M,x_0) \mid f_\#(\overline{\alpha}) = g_\#(\overline{\alpha})\}$ by $Coin(f_\#,g_\#)_{x_0}$, we have:

Corollary 3.5. *If* x_0 *is a coincidence of the pair* (f,g) *then the set* $p^{-1}(x_0)\cap Coin(\widetilde{f},\widetilde{g})$ *can be partitioned in* $\frac{\#Coin(\widetilde{f}_{*,x_0},\widetilde{g}_{*,x_0})}{\#j_{x_0}(Coin(f_\#,g_\#)_{x_0})}$ *disjoint subsets, each of then with* $\#j_{x_0}(Coin(f_\#,g_\#)_{x_0})$ *elements that are all Nielsen related (therefore contained in the same coincidence class of the pair* $(\widetilde{f},\widetilde{g})$*). Moreover, no two points of different subsets are Nielsen related.* □

Lemma 3.6. *Let* x_0, x_1 *be coincidence points in the same coincidence class of the pair* (f,g)*,* $\overline{\alpha}$ *be a path from* x_0 *to* x_1 *that realizes the Nielsen relation,* $\widetilde{x_0},\widetilde{x}'_0$ *be coincidence points of the pair* $(\widetilde{f},\widetilde{g})$ *such that* $p(\widetilde{x_0}) = p(\widetilde{x}'_0) = x_0$*, and* γ *the unique element of* $\mathfrak{D}(\widetilde{M})$ *such that* $\gamma(\widetilde{x_0}) = \widetilde{x}'_0$*. If* $\widetilde{\alpha}$ *and* $\widetilde{\alpha}'$ *are the two liftings of* $\overline{\alpha}$ *starting at* $\widetilde{x_0}$ *and* $\widetilde{x}'_0$ *respectively then:*

(i) $\widetilde{\alpha}(1)$ *and* $\widetilde{\alpha}'(1)$ *are coincidence points of the pair* $(\widetilde{f},\widetilde{g})$*;*
(ii) $\widetilde{\alpha}(1)$ *(*$\widetilde{\alpha}'(1)$*) is in the same coincidence class as* $\widetilde{x_0}$ *(*$\widetilde{x}'_0$*).*
(iii) $p(\widetilde{\alpha}(1)) = p(\widetilde{\alpha}'(1)) = x_1$*;*

(iv) $\gamma(\widetilde{\alpha}(1)) = \widetilde{\alpha}'(1)$.

(v) *If* $\overline{\alpha}$ *is a graph orientation-reversing-path (in this case* $x_0 R x_1$*) then* $\widetilde{\alpha}(1)$ *and* $\widetilde{\alpha}'(1)$ *are graph orientation-reverse-paths (in this case* $\widetilde{x}_0 R \widetilde{x}_1$ *and* $\widetilde{x}'_0 R \widetilde{x}'_1$*).*

Proof: (i), (ii) and (iii) are known (we can prove them using covering space theory). To prove (iv), we take $\overline{\gamma} \in \pi_1(M, x_0)$ such that $\mathfrak{D}(\widetilde{M}) \ni [\overline{\gamma}] = \gamma$ and we can see that projection of the path $\overline{\alpha} \circ \overline{\gamma} \circ \overline{\alpha}^{-1} \in \pi_1(M, x_1)$ in $\mathfrak{D}(\widetilde{M})$ coincides with γ.

To prove (v), we use [Dobreńko & Jezierski, lemma 2.1, page 77]. □

Theorem 3.7. *Let* M *and* N *be compact, closed manifolds of the same dimension,* $f, g : M \to N$ *be two maps such that* $Coin(f, g)$ *is finite, and* $p : \widetilde{M} \to M$ *and* $q : \widetilde{N} \to N$ *be finite coverings such that there exist lifts* $\widetilde{f}, \widetilde{g} : \widetilde{M} \to \widetilde{N}$ *of the pair* (f, g)*. If* $\widetilde{C}$ *is a coincidence class of the pair* $(\widetilde{f}, \widetilde{g})$ *then* $C = p(\widetilde{C})$ *is a coincidence class of the pair* (f, g) *and*

$$|ind|(\widetilde{f}, \widetilde{g}; \widetilde{C}) = \begin{cases} s \cdot k \ (mod\ 2) & \text{if } C \text{ is defective;} \\ s \cdot k & \text{otherwise,} \end{cases}$$

where $s = |ind|(f, g, C)$*,* $k = \#j(Coin(f_\#, g_\#)_{x_0})$ *and* $x_0 \in C$*.*

Proof: The fact that $C = p(\widetilde{C})$ is a coincidence class of the pair (f, g) is known. We choose a point $x_0 \in C$. Since $Coin(f, g)$ is finite we can suppose that a decomposition $C = \{x_1, x_2, \cdots, x_s; c_1, c'_1, c_2, c'_2, \cdots, c_n, c'_n\}$ is such that each x_i is free, and for all pairs c_j, c'_j we have $c_j R c'_j$.

Now we choose paths $\{\overline{\alpha}_i\}_i$, $2 \le i \le s$; $\{\overline{\beta}_j\}_j$ and $\{\overline{\gamma}_j\}_j$, $1 \le j \le n$ (see figure 1) such that:

- $\overline{\alpha}_i$ is a path in M from x_1 to x_i that realizes the Nielsen relation.
- $\overline{\beta}_j$ is a path in M from x_1 to c_j that realizes the Nielsen relation.
- $\overline{\gamma}_j$ is a graph-orientation-reversing path in M from c_j to c'_j.



FIGURE 1. The class C and the chosen paths.

For each element $\{x_1^k\}$ of $p^{-1}(x_1) \cap Coin(\widetilde{f}, \widetilde{g})$ (by Corollary 3.2 there were $\#Coin(\widetilde{f}_{*,x_1}, \widetilde{g}_{*,x_1})$ such elements) we take lifts $\{\widetilde{\alpha}_i^k\}_{i,k}$, $\{\widetilde{\beta}_j^k\}_{j,k}$ and $\{\widetilde{\phi}_j^k\}_{j,k}$ of the paths $\{\overline{\alpha}_i\}_i$, $\{\overline{\beta}_j\}_j$ and $\{\overline{\gamma}_j \circ \overline{\beta}_j\}_j$ respectively, starting at x_1^k.

Using Corollary 3.5 and Lemma 3.6 at the point $x_1 \in C$ we obtain that the set $p^{-1}(C) \cap Coin(\widetilde{f}, \widetilde{g})$ is the union of $\#Coin(\widetilde{f}_{*,x_1}, \widetilde{g}_{*,x_1})$ copies of the class C and these copies can be divided in $\frac{\#Coin(\widetilde{f}_{*,x_1},\widetilde{g}_{*,x_1})}{\#j_{x_1}(Coin(f_\#,g_\#)_{x_1})}$ disjoint coincidence classes of the pair $(\widetilde{f},\widetilde{g})$. Each one of these classes contains $\#j_{x_1}(Coin(f_\#,g_\#)_{x_1})$ copies of the class C.

The result follows by Corollary 3.4. □

4. Two folded orientable covering

Let M and N be manifolds of same dimension with M nonorientable and N orientable; $f,g : M \to N$ be two maps such that $Coin(f,g)$ is finite, and $p : \widetilde{M} \to M$ be the two-fold orientable covering of M. We define $\widetilde{f},\widetilde{g} : \widetilde{M} \to N$ by $\widetilde{f} = f \circ p$ and $\widetilde{g} = g \circ p$.

$$\begin{tikzcd}
\widetilde{M} \arrow[d, "p"'] \arrow[dr, "\widetilde{f}", "\widetilde{g}"'] & \\
M \arrow[r, "f", "g"'] & N
\end{tikzcd}$$

Lemma 4.1. *Under the above conditions, if C is a coincidence class of the pair (f,g) then $p^{-1}(C) \subset Coin(\widetilde{f},\widetilde{g})$ is such that:*

1. *$p^{-1}(C)$ can be divided in two disjoint sets $\widetilde{C}$ and $\widetilde{C}'$, of the same cardinality, such that $p(\widetilde{C}) = p(\widetilde{C}') = C$.*
2. *If $\widetilde{x_1},\widetilde{x_2} \in \widetilde{C}$ (or $\widetilde{C}'$) then $\widetilde{x_1}$ and $\widetilde{x_2}$ are in the same coincidence class of $(\widetilde{f},\widetilde{g})$*
3. *$\widetilde{C}$ and $\widetilde{C}'$ are in the same coincidence class of the pair $(\widetilde{f},\widetilde{g})$ if and only if C is defective.*

Proof: We need only to take $q : \widetilde{N} \to N$ as the identity map in the Corollary 3.2, Corollary 3.4 and Lemma 3.6. □

Corollary 4.2. *Under the hypotheses of lemma 4.1 we have:*

1. *If C is not defective then $\widetilde{C}$ and $\widetilde{C}'$ are two coincidence classes of the pair $(\widetilde{f},\widetilde{g})$ with $ind(\widetilde{f},\widetilde{g},\widetilde{C}) = -ind(\widetilde{f},\widetilde{g},\widetilde{C}')$ and $|ind|(\widetilde{f},\widetilde{g},\widetilde{C}) = |ind(f,g,C)|$.*
2. *If C is defective then $\widetilde{C} \cup \widetilde{C}'$ is a unique coincidence class of the pair $(\widetilde{f},\widetilde{g})$ with $ind(\widetilde{f},\widetilde{g},\widetilde{C} \cup \widetilde{C}') = 0$.*

Proof: The first part is a consequence of lemma 4.1, and the second can be proved using the fact that $\widetilde{M}$ is the double orientable covering of M. It is useful to remember that the pair $(\widetilde{f},\widetilde{g})$ is a pair of maps between orientable manifolds. □

Corollary 4.3. *Under de hypotheses of lemma 4.1 we have:*

1. *$L(\widetilde{f},\widetilde{g}) = 0$;*
2. *$N(\widetilde{f},\widetilde{g})$ is even;*
3. *$N(f,g) \geq \frac{N(\widetilde{f},\widetilde{g})}{2}$;*

(4) *If* $N(\widetilde{f},\widetilde{g}) = 0$ *then all coincidence classes with nonzero semi-index of the pair* (f,g) *are defective.*

Proof: We need only to see that $p(Coin(\widetilde{f},\widetilde{g})) = Coin(f,g)$, and that in the pair $(\widetilde{f},\widetilde{g})$ we "lose" the defective classes. □

5. Applications

Theorem 5.1. *Let* $f, g : M \to N$ *be two maps between compact closed manifolds of the same dimension such that* M *is nonorientable and* N *is orientable. Suppose that* N *is such that for all orientable manifolds* M' *of the same dimension of* N *and all pairs of maps* $f', g' : M' \to N$ *we have* $L(f',g') = 0 \Rightarrow N(f',g') = 0$. *Then all coincidence classes with nonzero semi-index of the pair* (f,g) *are defective.*

Proof: The hypotheses on N are enough to show, using the notation of the proof of theorem 4.1, that $N(\widetilde{f},\widetilde{g}) = 0$. So by corollary 4.3, all coincidence classes with nonzero semi-index of the pair (f,g) are defective. □

We note that the hypotheses on the manifold N in theorem 5.1, in dimension greater then 2, are equivalent to the converse of Lefschetz theorem. In dimension 2 these hypotheses are not equivalent but necessary for the converse of Lefschetz theorem.

Remark 5.2. *The following manifolds satisfy the hypotheses on the manifold* N *in the theorem 5.1:*

1. *Jiang spaces (*[Gonçalves & Wong 1, Corollary 1]*).*
2. *Nilmanifolds (*[Gonçalves & Wong 2, Theorem 5]*).*
3. *Homogeneous spaces of a compact connected Lie group* G *by a finite subgroup* K *(*[Gonçalves & Wong 1, Theorem 4]*).*
4. *Suitable manifolds (*[Wong, Theorem 3 e Theorem 4]*) .*

Acknowledgements

This work was made during a postdoctoral year of the author at Laboratoire Émile Picard - Université Paul Sabatier (Toulouse - France). We would like to thank John Guaschi and Claude Hayat-Legrand for the invitation and hospitality, and Peter N.-S. Wong for helpful conversations. This work was supported by Capes-BEX0755/02-8 (international cooperation Capes-Cofecub project number 364/01).

References

[Brown & Schirmer] Brown, Robert F.; Schirmer, Helga. *Nielsen root theory and Hopf degree theory*, Pacific J. Math., 198 (2001), No 1, 49–80.

[Dobreńko & Jezierski] Dobreńko, R.; Jezierski, J. *The coincidence Nielsen theory on non-orientable manifolds*, Rocky Mountain J. math. 23 (1993), 67–85.

[Gonçalves & Jezierski] Gonçalves, Daciberg L.; Jezierski, J. *Lefschetz coincidence formula on non-orientable manifolds*, Fund. Math. 153 (1997), 01–23.

[Gonçalves & Wong 1] Gonçalves, Daciberg L.; Wong, Peter N.-S. *Homogeneous spaces in coincidence theory*, In: *Proceedings of the X Encontro Brasileiro de Topologia (São Carlos, Brasil)*, (P. Schweitzer, ed.), Sociedade Brasileira de Matemática, Matemática Contemporânea, vol 13 (1997), 143–158.

[Gonçalves & Wong 2] Gonçalves, Daciberg L.; Wong, Peter N.-S. *Nilmanifolds are Jiang-type spaces for coincidences*, Forum Math. 13 (2001), 133–141.

[Jezierski 1] Jezierski, J. *The semi index product formula*, Fund. Math. 140 (1992), 99–120.

[Jezierski 2] Jezierski, J. *The Nielsen coincidence theory on topological manifolds*, Fund. Math. 143 (1993), 167–178.

[Olum] Olum, P. *Mappings of manifolds and the notion of degree*, Ann. Math. 58 (1953), 458–480.

[Skora] Skora, R. *The degree of a map between surfaces*, Ann. Math. 276 (1987), 415–423.

[Wong] Wong, Peter N.-S. *Coincidences of maps into homogeneous spaces*, Manuscripta Math. 98 (1998), 243–254.

E-mail address: `vendrus@picard.ups-tlse.fr`